


PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

J. Wei

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C11D 3/386, 3/16		A1	(11) International Publication Number: WO 96/41859
			(43) International Publication Date: 27 December 1996 (27.12.96)
(21) International Application Number: PCT/DK96/00252		(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 10 June 1996 (10.06.96)		Published With international search report.	
(30) Priority Data: 0674/95 13 June 1995 (13.06.95) DK 0983/95 7 September 1995 (07.09.95) DK			
(71) Applicant (for all designated States except US): NOVO NORDISK A/S [DK/DK]; Novo Allé, DK-2880 Bagsværd (DK).			
(72) Inventors; and (75) Inventors/Applicants (for US only): NIELSEN, Lone, Kirsten [DK/DK]; Novo Nordisk a/s, Novo Allé, DK-2880 Bagsværd (DK). DEANE-WRAY, Allison [GB/GB]; 7 Bridge Meadows, Liss, Hampshire GU33 7JY (GB).			
(74) Agent: NOVO NORDISK A/S; Corporate Patents, Novo Allé, DK-2880 Bagsværd (DK).			
(54) Title: 4-SUBSTITUTED-PHENYL-BORONIC ACIDS AS ENZYME STABILIZERS			
(57) Abstract			
<p>This invention relates to a liquid composition comprising an enzyme and a phenyl boronic acid derivative enzyme stabilizer of formula (I), wherein R is selected from the group consisting of hydrogen, hydroxy, C₁-C₆ alkyl, substituted C₁-C₆ alkyl, C₁-C₆ alkenyl and substituted C₁-C₆ alkenyl.</p>		 <p style="text-align: right;">(I)</p>	

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

4-SUBSTITUTED-PHENYL-BORONIC ACIDS AS ENZYME STABILIZERS

FIELD OF INVENTION

5 This invention relates to a liquid composition, in particular to a liquid detergent composition, comprising an enzyme and an improved enzyme stabilizer.

BACKGROUND OF THE INVENTION

10

Storage stability problems are well known with liquids containing enzyme(s). Especially in enzyme-containing liquid detergents a major problem, in particular if the detergent contains protease, is that of ensuring enzyme activity over time.

The prior art has dealt extensively with improving the storage stability, for example by adding a protease inhibitor.

Boric acid and boronic acids are known to reversibly inhibit proteolytic enzymes. A discussion of the inhibition of one serine protease, subtilisin, by boronic acid is provided in Molecular & Cellular Biochemistry 51, 1983, pp. 5-32.

Boronic acids have very different capacities as subtilisin inhibitors. Boronic acids containing only alkyl groups such as methyl, butyl or 2-cyclohexylethyl are poor inhibitors with methylboronic acid as the poorest inhibitor, whereas boronic acids bearing aromatic groups such as phenyl, 4-methoxyphenyl or 3,5-dichlorophenyl are good inhibitors with 3,5-dichlorophenylboronic acid as a particularly effective one (see Keller et al, Biochem. Biophys. Res. Com. 176, 1991, pp. 401-405).

It is also claimed that aryl boronic acids which have a substitution at the 3-position relative to boron are unexpectedly good reversible protease inhibitors. Especially,

acetamidophenyl boronic acid is claimed to be a superior inhibitor of proteolytic enzymes (see WO 92/19707).

The inhibition constant (K_i) is ordinarily used as a measure of capacity to inhibit enzyme activity, with a low K_i indicating a more potent inhibitor. However, it has earlier been found that the K_i values of boronic acids do not always tell how effective inhibitors are (see for instance WO 92/19707).


10

SUMMARY OF THE INVENTION

In this invention it is surprisingly found that phenyl boronic acid derivatives substituted in the para-position with a $>C=O$ adjacent to the phenyl boronic acid have extraordinary good capacities as enzyme stabilizers in liquids.

Accordingly, the present invention relates to a liquid composition comprising an enzyme and a phenyl boronic acid derivative enzyme stabilizer of the following formula:

20




wherein R is selected from the group consisting of hydrogen, hydroxy, C_1 - C_6 alkyl, substituted C_1 - C_6 alkyl, C_1 - C_6 alkenyl and substituted C_1 - C_6 alkenyl.

DETAILED DISCLOSURE OF THE INVENTION

30

One embodiment of the present invention provides a liquid composition comprising an enzyme and a phenyl boronic acid derivative enzyme stabilizer of the following formula:



wherein R is selected from the group consisting of hydrogen, 5 hydroxy, C₁-C₆ alkyl, substituted C₁-C₆ alkyl, C₁-C₆ alkenyl and substituted C₁-C₆ alkenyl.

A preferred embodiment of the present invention provides a liquid composition comprising an enzyme and a phenyl boronic acid derivative enzyme stabilizer of the 10 formula disclosed above, wherein R is a C₁-C₆ alkyl, in particular wherein R is CH₃, CH₃CH₂ or CH₃CH₂CH₂, or wherein R is hydrogen.

A further preferred embodiment of the present invention provides a liquid detergent composition comprising a 15 surfactant, an enzyme and a phenyl boronic acid derivative enzyme stabilizer of the formula disclosed above.

Preparation of Phenyl Boronic Acid Derivatives

Phenyl boronic acid derivatives may be prepared 20 using methods well known to those skilled in the art, for example by using a Grignard preparation:

The Grignard reagent is prepared by the slow dropwise addition of the appropriate bromobenzene starting material in anhydrous ether to magnesium turnings in anhydrous 25 ether. The anhydrous ether may be, e.g., sodium dried diethylether or sodium dried tetrahydrofuran. The reaction is encouraged by the addition of a small iodine crystal.

Trimethylborate or tri-n-butylborate in anhydrous ether (e.g. sodium dried diethylether or sodium dried 30 tetrahydrofuran) is cooled to about -70°C and the Grignard reagent is added dropwise over a period of approximately 2 hours while keeping the borate solution at about -70°C and continuously agitating.

The reaction mixture is allowed to warm to room temperature overnight whereupon it is hydrolysed by the dropwise addition of cold dilute sulphuric acid. The ether layer is separated and the aqueous layer extracted with ether. 5 The ether containing fractions are combined and the solvent removed. The residue is made distinctly alkaline and any methanol or butanol so formed is removed. The alkaline solution is made acidic and cooled and the resulting crystals of desired boronic acid are removed by filtration. All 10 products are preferably recrystallized from distilled water or some other appropriate solvent.

Preparation of, e.g., 4-formyl-phenyl-boronic acid, using the method disclosed above, has been described in Chem. Ber. 123, 1990, pp. 1841-1843.

15 The phenyl boronic acids may also be prepared using either direct lithiation of the benzene and/or lithiation of the bromide.

Any nuclear substitution or protection of functional groups may be achieved by using standard methods well known to 20 those skilled in the art.

Stabilizers

According to the invention the liquid composition may contain up to 500 mM of the stabilizer (the phenyl boronic 25 acid derivative), preferably the detergent composition may contain 0.001-250 mM of the stabilizer, more preferably the liquid composition may contain 0.005-100 mM of the stabilizer, most preferably the liquid composition may contain 0.01-10 mM of the stabilizer. The phenyl boronic acid derivative may be 30 an acid or the alkali metal salt of said acid.

Enzymes

According to the invention the liquid composition contains at least one enzyme. The enzyme may be any 35 commercially available enzyme, in particular an enzyme

selected from the group consisting of proteases, amylases, lipases, cellulases, oxidoreductases and any mixture thereof. Mixtures of enzymes from the same class (e.g. proteases) are also included.

protease preferred

5 According to the invention a liquid composition comprising a protease is preferred; more preferred is a liquid composition comprising two or more enzymes in which the first enzyme is a protease and the second enzyme is selected from the group consisting of amylases, lipases, cellulases and
10 oxidoreductases; even more preferred is a liquid composition in which the first enzyme is a protease and the second enzyme is a lipase.

The amount of enzyme used in the liquid composition varies according to the type of enzyme(s). The amount of each
15 enzyme will typically be 0.04-40 μ M, in particular 0.2-30 μ M, especially 0.4-20 μ M (generally 1-1000 mg/l, in particular 5-750 mg/l, especially 10-500 mg/l) calculated as pure enzyme protein.

Proteases: Suitable proteases include those of animal, 20 vegetable or microbial origin. Microbial origin is preferred. Chemically or genetically modified mutants are included. The protease may be a serine protease, preferably an alkaline microbial protease or a trypsin-like protease. Examples of alkaline proteases are subtilisins, especially those derived
25 from Bacillus, e.g., subtilisin Novo, subtilisin Carlsberg, subtilisin 309, subtilisin 147 and subtilisin 168 (described in WO 89/06279). Examples of trypsin-like proteases are trypsin (e.g. of porcine or bovine origin) and the Fusarium protease described in WO 89/06270.

30 Preferred commercially available protease enzymes include those sold under the tradenames Alcalase, Savinase, Primase, Durazym, and Esperase by Novo Nordisk A/S (Denmark), those sold under the tradename Maxatase, Maxacal, Maxapem and Properase by Gist-Brocades, those sold under the tradename
35 Purafect and Purafect OXP by Genencor International, and

those sold under the tradename Opticlean and Optimase by Solvay Enzymes.

Lipases: Suitable lipases include those of bacterial or fungal origin. Chemically or genetically modified mutants are 5 included.

Examples of useful lipases include a Humicola lanuginosa lipase, e.g., as described in EP 258 068 and EP 305 216, a Rhizomucor miehei lipase, e.g., as described in EP 238 023, a Candida lipase, such as a C. antarctica lipase, e.g., the 10 C. antarctica lipase A or B described in EP 214 761, a Pseudomonas lipase such as a P. pseudoalcaligenes and P. alcaligenes lipase, e.g., as described in EP 218 272, a P. cepacia lipase, e.g., as described in EP 331 376, a P. stutzeri lipase, e.g., as disclosed in BP 1,372,034, a P. fluorescens 15 lipase, a Bacillus lipase, e.g., a B. subtilis lipase (Dartois et al., (1993), Biochemica et Biophysica acta 1131, 253-260), a B. stearothermophilus lipase (JP 64/744992) and a B. pumilus lipase (WO 91/16422).

Furthermore, a number of cloned lipases may be useful, 20 including the Penicillium camembertii lipase described by Yamaguchi et al., (1991), Gene 103, 61-67), the Geotricum candidum lipase (Schimada, Y. et al., (1989), J. Biochem. 106, 383-388), and various Rhizopus lipases such as a R. delemar lipase (Hass, M.J et al., (1991), Gene 109, 117-113), a R. 25 niveus lipase (Kugimiya et al., (1992), Biosci. Biotech. Biochem. 56, 716-719) and a R. oryzae lipase.

Other types of lipolytic enzymes such as cutinases may also be useful, e.g., a cutinase derived from Pseudomonas mendocina as described in WO 88/09367, or a cutinase derived 30 from Fusarium solani pisi (e.g. described in WO 90/09446).

Especially suitable lipases are lipases such as M1 Lipase™, Luma fast™ and Lipomax™ (Genencor), Lipolase™ and Lipolase Ultra™ (Novo Nordisk A/S), and Lipase P "Amano" (Amano Pharmaceutical Co. Ltd.).

Amylases: Suitable amylases (α and/or β) include those of bacterial or fungal origin. Chemically or genetically modified mutants are included. Amylases include, for example, α -amylases obtained from a special strain of B. licheniformis,
5 described in more detail in British Patent Specification No. 1,296,839. Commercially available amylases are Duramyl™, Termamyl™, Fungamyl™ and BAN™ (available from Novo Nordisk A/S) and Rapidase™ and Maxamyl P™ (available from Gist-Brocades).

Cellulases: Suitable cellulases include those of bacterial or fungal origin. Chemically or genetically modified mutants are included. Suitable cellulases are disclosed in US
10 4,435,307, which discloses fungal cellulases produced from Humicola insolens. Especially suitable cellulases are the cellulases having color care benefits. Examples of such cel-
15 lulases are cellulases described in European patent application No. 0 495 257.

Commercially available cellulases is Celluzyme™ produced by a strain of Humicola insolens, (Novo Nordisk A/S), and KAC-500(B)™ (Kao Corporation).

20 Oxidoreductases: Any oxidoreductase suitable for use in a liquid composition, e.g., peroxidases or oxidases such as laccases, can be used herein. Suitable peroxidases herein include those of plant, bacterial or fungal origin. Chemically or genetically modified mutants are included. Examples of
25 suitable peroxidases are those derived from a strain of Coprinus, e.g., C. cinerius or C. macrorrhizus, or from a strain of Bacillus, e.g., B. pumilus, particularly peroxidase according to WO 91/05858. Suitable laccases herein include those of bacterial or fungal origin. Chemically or genetically
30 modified mutants are included. Examples of suitable laccases are those obtainable from a strain of Trametes, e.g., T. villosa or T. versicolor, or from a strain of Coprinus, e.g., C. cinereus, or from a strain of Myceliophthora, e.g., M. thermophila.

35

Detergents

According to the invention the liquid detergent composition will beside enzyme(s) and stabilizer comprise a surfactant. The detergent composition may, e.g., be a laundry
5 detergent composition or a dishwashing detergent composition.

The detergent may be aqueous, typically containing up to 70 % water and 0-30 % organic solvent, or nonaqueous.

The detergent composition comprises one or more surfactants, each of which may be anionic, nonionic, cationic,
10 or amphoteric (zwitterionic). The detergent will usually contain 0-50% of anionic surfactant such as linear alkylbenzenesulfonate (LAS), alpha-olefinsulfonate (AOS), alkyl sulfate (fatty alcohol sulfate) (AS), alcohol ethoxysulfate (AEOS or AES), secondary alkanesulfonates (SAS), alpha-sulfo
15 fatty acid methyl esters, alkyl- or alkenylsuccinic acid, or soap. It may also contain 0-40% of nonionic surfactant such as alcohol ethoxylate (AEO or AE), alcohol propoxylate, carboxylated alcohol ethoxylates, nonylphenol ethoxylate, alkylpolyglycoside, alkyldimethylamine oxide, ethoxylated
20 fatty acid monoethanolamide, fatty acid monoethanolamide, or polyhydroxy alkyl fatty acid amide (e.g. as described in WO 92/06154).

Normally the detergent contains 1-65% of a detergent builder, but some dishwashing detergents may contain even up
25 to 90% of a detergent builder, or complexing agent such as zeolite, diphosphate, triphosphate, phosphonate, citrate, nitrilotriacetic acid (NTA), ethylenediaminetetraacetic acid (EDTA), diethylenetriaminepentaacetic acid (DTMPA), alkyl- or alkenylsuccinic acid, soluble silicates or layered silicates
30 (e.g. SKS-6 from Hoechst).

The detergent builders may be subdivided into phosphorus-containing and non-phosphorous-containing types. Examples of phosphorus-containing inorganic alkaline detergent builders include the water-soluble salts, especially alkali
35 metal pyrophosphates, orthophosphates, polyphosphates and

phosphonates. Examples of non-phosphorus-containing inorganic builders include water-soluble alkali metal carbonates, borates and silicates as well as layered disilicates and the various types of water-insoluble crystalline or amorphous alumino silicates of which zeolites is the best known representative.

Examples of suitable organic builders include alkali metal, ammonium or substituted ammonium salts of succinates, malonates, fatty acid malonates, fatty acid sulphonates, carboxymethoxy succinates, polyacetates, carboxylates, polycarboxylates, aminopolycarboxylates and polyacetyl carboxylates. The detergent may also be unbuild, i.e. essentially free of detergent builder.

The detergent may comprise one or more polymers. Examples are carboxymethylcellulose (CMC), poly(vinylpyrrolidone) (PVP), polyethyleneglycol (PEG), poly(vinyl alcohol) (PVA), polycarboxylates such as polyacrylates, polymaleates, maleic/acrylic acid copolymers and lauryl methacrylate/acrylic acid copolymers.

The detergent composition may contain bleaching agents of the chlorine/bromine-type or the oxygen-type. The bleaching agents may be coated or encapsulated. Examples of inorganic chlorine/bromine-type bleaches are lithium, sodium or calcium hypochlorite or hypobromite as well as chlorinated trisodium phosphate. The bleaching system may also comprise a H_2O_2 source such as perborate or percarbonate which may be combined with a peracid-forming bleach activator such as tetraacetylenediamine (TAED) or nonanoyloxybenzene-sulfonate (NOBS).

Examples of organic chlorine/bromine-type bleaches are heterocyclic N-bromo and N-chloro imides such as trichloroisocyanuric, tribromoisocyanuric, dibromoisocyanuric and dichloroisocyanuric acids, and salts thereof with water solubilizing cations such as potassium and sodium. Hydantoin compounds are also suitable. The bleaching system may also

comprise peroxyacids of, e.g., the amide, imide, or sulfone type.

In dishwashing detergents the oxygen bleaches are preferred, for example in the form of an inorganic persalt, preferably with a bleach precursor or as a peroxy acid compound. Typical examples of suitable peroxy bleach compounds are alkali metal perborates, both tetrahydrates and monohydrates, alkali metal percarbonates, persilicates and perphosphates. Preferred activator materials are TAED or NOBS.

The enzyme(s) of the detergent composition of the invention may additionally be stabilized using conventional stabilizing agents, e.g., a polyol such as propylene glycol or glycerol, a sugar or sugar alcohol, or lactic acid.

The detergent may also contain other conventional detergent ingredients such as, e.g., fabric conditioners including clays, deflocculant material, foam boosters/foam depressors (in dishwashing detergents foam depressors), suds suppressors, anti-corrosion agents, soil-suspending agents, anti-soil-redeposition agents, dyes, dehydrating agents, bactericides, optical brighteners, or perfume.

The pH (measured in aqueous solution at use concentration) will usually be neutral or alkaline, e.g. in the range of 7-11.

Particular forms of laundry detergent compositions within the scope of the invention include:

1) An aqueous liquid detergent composition comprising

Linear alkylbenzenesulfonate (calculated as acid)	15	- 21%
Alcohol ethoxylate (e.g. C ₁₂₋₁₅ alcohol, 7 EO or C ₁₂₋₁₅ alcohol, 5 EO)	12	- 18%
Soap as fatty acid (e.g. oleic acid)	3	- 13%
Alkenylsuccinic acid (C ₁₂₋₁₄)	0	- 13%
Aminoethanol	8	- 18%
Citric acid	2	- 8%

Phosphonate	0 - 3%
Polymers (e.g. PVP, PEG)	0 - 3%
Borate (as B ₄ O ₇)	0 - 2%
Ethanol	0 - 3%
Propylene glycol	8 - 14%
Enzymes (calculated as pure enzyme protein)	0.0001 - 0.1%
Minor ingredients (e.g. dispersants, suds suppressors, perfume, optical brightener)	0 - 5%

2) An aqueous structured liquid detergent composition comprising

Linear alkylbenzenesulfonate (calculated as acid)	15 - 21%
Alcohol ethoxylate (e.g. C ₁₂₋₁₅ alcohol, 7 EO, or C ₁₂₋₁₅ alcohol, 5 EO)	3 - 9%
Soap as fatty acid (e.g. oleic acid)	3 - 10%
Zeolite (as NaAlSiO ₄)	14 - 22%
Potassium citrate	9 - 18%
Borate (as B ₄ O ₇)	0 - 2%
Carboxymethylcellulose	0 - 2%
Polymers (e.g. PEG, PVP)	0 - 3%
Anchoring polymers such as, e.g., lauryl methacrylate/acrylic acid copolymer; molar ratio 25:1; MW 3800	0 - 3%
Glycerol	0 - 5%
Enzymes (calculated as pure enzyme protein)	0.0001 - 0.1%
Minor ingredients (e.g. dispersants, suds suppressors, perfume, optical brighteners)	0 - 5%

5 3) An aqueous liquid detergent composition comprising

Linear alkylbenzenesulfonate	
------------------------------	--

(calculated as acid)	15	- 23%
Alcohol ethoxysulfate (e.g. C ₁₂₋₁₅ alcohol, 2-3 EO)	8	- 15%
Alcohol ethoxylate (e.g. C ₁₂₋₁₅ alcohol, 7 EO, or C ₁₂₋₁₅ alcohol, 5 EO)	3	- 9%
Soap as fatty acid (e.g. lauric acid)	0	- 3%
Aminoethanol	1	- 5%
Sodium citrate	5	- 10%
Hydrotrope (e.g. sodium toluensulfonate)	2	- 6%
Borate (as B ₄ O ₇)	0	- 2%
Carboxymethylcellulose	0	- 1%
Ethanol	1	- 3%
Propylene glycol	2	- 5%
Enzymes (calculated as pure enzyme protein)	0.0001	- 0.1%
Minor ingredients (e.g. polymers, dispersants, perfume, optical brighteners)	0	- 5%

4) An aqueous liquid detergent composition comprising

Linear alkylbenzenesulfonate (calculated as acid)	20	- 32%
Alcohol ethoxylate (e.g. C ₁₂₋₁₅ alcohol, 7 EO, or C ₁₂₋₁₅ alcohol, 5 EO)	6	- 12%
Aminoethanol	2	- 6%
Citric acid	8	- 14%
Borate (as B ₄ O ₇)	1	- 3%
Polymer (e.g. maleic/acrylic acid copolymer, anchoring polymer such as, e.g., lauryl methacrylate/acrylic acid copolymer)	0	- 3%
Glycerol	3	- 8%
Enzymes (calculated as pure enzyme protein)	0.0001	- 0.1%

Minor ingredients (e.g. hydrotropes, dispersants, perfume, optical brighteners)	0 - 5%
---	--------

5) Detergent formulations as described in 1) - 4) wherein all or part of the linear alkylbenzenesulfonate is replaced by (C₁₂-C₁₈) alkyl sulfate.

5

6) Detergent formulations as described in 1) - 5) which contain a stabilized or encapsulated peracid, either as an additional component or as a substitute for already specified bleach systems.

10 7) Detergent composition formulated as a nonaqueous detergent liquid comprising a liquid nonionic surfactant such as, e.g., linear alkoxyated primary alcohol, a builder system (e.g. phosphate), enzyme and alkali. The detergent may also comprise anionic surfactant and/or a bleach system.

15 Particular forms of dishwashing detergent compositions within the scope of the invention include:

1) LIQUID DISHWASHING COMPOSITION WITH CLEANING SURFACTANT SYSTEM

20

Nonionic surfactant	0 - 1.5%
Octadecyl dimethylamine N-oxide dihydrate	0 - 5%
80:20 wt.C18/C16 blend of octadecyl dimethylamine N-oxide dihydrate and hexadecyldimethyl amine N-oxide dihydrate	0 - 4%
70:30 wt.C18/C16 blend of octadecyl bis (hydroxyethyl)amine N-oxide anhydrous and hexadecyl bis (hydroxyethyl)amine N-oxide anhydrous	0 - 5%
C ₁₃ -C ₁₅ alkyl ethoxysulfate with an average degree of ethoxylation of 3	0 - 10%
C ₁₂ -C ₁₅ alkyl ethoxysulfate with an average degree of ethoxylation of 3	0 - 5%

C ₁₂ -C ₁₅ ethoxylated alcohol with an average degree of ethoxylation of 12	0	- 5%
A blend of C ₁₂ -C ₁₅ ethoxylated alcohols with an average degree of ethoxylation of 9	0	- 6.5%
A blend of C ₁₂ -C ₁₅ ethoxylated alcohols with an average degree of ethoxylation of 30	0	- 4%
Sodium disilicate	0	- 33%
Sodium tripolyphosphate	0	- 46%
Sodium citrate	0	- 28%
Citric acid	0	- 29%
Sodium carbonate	0	- 20%
Sodium perborate monohydrate	0	- 11.5%
Tetraacetythylenediamine (TAED)	0	- 4%
Maleic acid/acrylic acid copolymer	0	- 7.5%
Sodium sulphate	0	- 12.5%
Enzymes	0.0001	- 0.1%

2) NON-AQUEOUS LIQUID AUTOMATIC DISHWASHING COMPOSITION

Liquid nonionic surfactant (e.g. alcohol ethoxylates)	2.0	- 10.0%
Alkali metal silicate	3.0	- 15.0%
Alkali metal phosphate	20.0	- 40.0%
Liquid carrier selected from higher glycols, polyglycols, polyoxides, glycolethers	25.0	- 45.0%
Stabilizer (e.g. a partial ester of phosphoric acid and a C ₁₆ -C ₁₈ alkanol)	0.5	- 7.0%
Foam suppressor (e.g. silicone)	0	- 1.5%
Enzymes	0.0001	- 0.1%

5 3) NON-AQUEOUS LIQUID DISHWASHING COMPOSITION

Liquid nonionic surfactant (e.g. alcohol ethoxylates)	2.0	- 10.0%
Sodium silicate	3.0	- 15.0%

Alkali metal carbonate	7.0 - 20.0%
Sodium citrate	0.0 - 1.5%
Stabilizing system (e.g. mixtures of finely divided silicone and low molecular weight dialkyl polyglycol ethers)	0.5 - 7.0%
Low molecule weight polyacrylate polymer	5.0 - 15.0%
Clay gel thickener (e.g. bentonite)	0.0 - 10.0%
Hydroxypropyl cellulose polymer	0.0 - 0.6%
Enzymes	0.0001 - 0.1%
Liquid carrier selected from higher lycols, polyglycols, polyoxides and glycol ethers	Balance

4) THIXOTROPIC LIQUID AUTOMATIC DISHWASHING COMPOSITION

C ₁₂ -C ₁₄ fatty acid	0 - 0.5%
Block co-polymer surfactant	1.5 - 15.0%
Sodium citrate	0 - 12%
Sodium triphosphate	0 - 15%
Sodium carbonate	0 - 8%
Aluminium tristearate	0 - 0.1%
Sodium cumene sulphonate	0 - 1.7%
Polyacrylate thickener	1.32 - 2.5%
Sodium polyacrylate	2.4 - 6.0%
Boric acid	0 - 4.0%
Sodium formate	0 - 0.45%
Calcium formate	0 - 0.2%
Sodium n-decydiphenyl oxide disulphonate	0 - 4.0%
Monoethanol amine (MEA)	0 - 1.86%
Sodium hydroxide (50%)	1.9 - 9.3%
1,2-Propanediol	0 - 9.4%
Enzymes	0.0001 - 0.1%
Suds suppressor, dye, perfumes,	

water	Balance
-------	---------

5) LIQUID AUTOMATIC DISHWASHING COMPOSITION

Alcohol ethoxylate	0	- 20%
Fatty acid ester sulphonate	0	- 30%
Sodium dodecyl sulphate	0	- 20%
Alkyl polyglycoside	0	- 21%
Oleic acid	0	- 10%
Sodium disilicate monohydrate	18	- 33%
Sodium citrate dihydrate	18	- 33%
Sodium stearate	0	- 2.5%
Sodium perborate monohydrate	0	- 13%
Tetraacetylenediamine (TAED)	0	- 8%
Maleic acid/acrylic acid copolymer	4	- 8%
Enzymes	0.0001	- 0.1%

5 6) LIQUID AUTOMATIC DISHWASHING COMPOSITION CONTAINING PROTECTED BLEACH PARTICLES

Sodium silicate	5	- 10%
Tetrapotassium pyrophosphate	15	- 25%
Sodium triphosphate	0	- 2%
Potassium carbonate	4	- 8%
Protected bleach particles, e.g. chlorine	5	- 10%
Polymeric thickener	0.7	- 1.5%
Potassium hydroxide	0	- 2%
Enzymes	0.0001	- 0.1%
Water	Balance	

7) Automatic dishwashing compositions as described in 1) and 10 5), wherein perborate is replaced by percarbonate.

8) Automatic dishwashing compositions as described in 1), which additionally contain a manganese catalyst. The manganese

catalyst may, e.g., be one of the compounds described in "Efficient manganese catalysts for low-temperature bleaching", Nature 369, 1994, pp. 637-639.

5 Tests of Stabilizers

According to the invention the effectiveness of each stabilizer may be tested in one or more of the following tests:

10 a) Storage Stability Test in Liquid Detergent: Enzyme(s) and stabilizer are added to a liquid detergent formulation and stored at well defined conditions. The enzyme activity of each enzyme is determined as a function of time, e.g. after 0, 3, 7 and 14 days.

15 To calculate the inhibition efficiency from the storage stability data a reaction mechanism is proposed. The following reactions give a relatively simple, but yet plausible, mechanism for a liquid detergent containing protease (P), lipase (L), and inhibitor (I):


20

I) Autodigestion of protease:




II) Denaturation of protease:


25



III) Inhibition of protease:



30 IV) Protease digestion of inhibited enzyme:




V) Denaturation of inhibited enzyme:



35

VI) Protease digestion of lipase:



VII) Denaturation of lipase:



where D_P and D_L are denatured (i.e. non-active) protease and lipase.

From these reactions three coupled differential
10 equations are derived describing the deactivation of P, L and
PI. The reaction rate constants are derived from storage
stability data by the use of a parameter estimation method
(Gauss-Newton with the Levenberg modification). The storage
stability data give the concentration of (P+PI) and L as a
15 function of time.

Reaction III is much faster than the other reactions
and equilibrium is assumed in the calculations. Reaction IV is
excluded from the system to reduce the number of parameters
thereby describing the stability of the inhibited enzyme by
20 only one reaction rate constant (from equation V).

In all experiments there is a large surplus of
inhibitor molecules compared to protease molecules, i.e. a
constant concentration of inhibitor (corresponding to the
added amount of inhibitor) is a reasonable assumption.

25 The specific values of the reaction rate constants
are somewhat sensitive to small variations in the data, but
the sensitivity is reduced significantly by giving the results
relatively to the value from Boric Acid. An improvement factor
is thus derived:

30

$$IF_I = \frac{K_I(\text{Boric Acid})}{K_I(\text{Inhibitor})}$$

35 IF_I measures the inhibition efficiency given by the inhibition
constants K_I from reaction III.

b) Determination of K_i : The inhibition constant K_i may be determined by using standard methods, for reference see Keller et al, Biochem. Biophys. Res. Com. 176, 1991, pp.401-405; J. 5 Bieth in Bayer-Symposium "Proteinase Inhibitors", pp. 463-469, Springer-Verlag, 1974 and Lone Kierstein Hansen in "Determination of Specific Activities of Selected Detergent Proteases using Protease Activity, Molecular Weights, Kinetic Parameters and Inhibition Kinetics", PhD-report, Novo Nordisk 10 A/S and University of Copenhagen, 1991.

The invention is further illustrated in the following examples which are not intended to be in any way limiting to the scope of the invention as claimed.

15

EXAMPLE 1Preparation of 4-Formyl-Phenyl-Boronic Acid

20 4-Formyl-phenyl-boronic acid may be prepared as disclosed in Chem. Ber. 123, 1990, pp. 1841-1843, or it may be bought at Lancaster Synthesis GmbH (4-Formylbenzeneboronic acid).

EXAMPLE 2

25

Determination of K_i

The inhibition constant K_i for the inhibition of Savinase™ (available from Novo Nordisk A/S) was determined using standard methods under the following conditions:

30

Substrate: Succinyl-Alanine-Alanine-Proline-Phenylalanine-para-nitro-anilide = SAAPFpNA (Sigma S-7388).

Buffer: 0.1 M Tris-HCl pH 8.6; 25°C.

35

Enzyme concentration in assay:Savinase: 1×10^{-10} - 3×10^{-10} M

The initial rate of substrate hydrolysis was determined at nine substrate concentrations in the range of 0.01 to 2 mM using a Cobas Fara automated spectrophotometer. The kinetic parameters V_{max} and K_m were determined using ENZFITTER (a non-linear regression data analysis program).

k_{cat} was calculated from the equation $V_{max} = k_{cat} \times [E_0]$. The concentration of active enzyme $[E_0]$ was determined by active site titration using tight-binding protein proteinase inhibitors. The inhibition constant K_i was calculated from plots of K_m/k_{cat} as a function of the concentration of inhibitor. The inhibitors were assumed to be 100% pure and the molar concentrations were determined using weighing numbers and molecular weights.

The results of the inhibition constants K_i of the phenyl boronic acid derivative enzyme stabilizers tested are listed below:

20

Inhibitor:	K_i (Savinase):

Boric acid	20 mM
4-formyl-phenyl-boronic acid	0.3 mM

25

For comparison reasons acetamidophenyl boronic acid was also tested in the same system giving the following results:

Inhibitor:	K_i (Savinase):

Boric acid	20 mM
acetamidophenyl boronic acid	1 mM

35

It appears from the results given above that the inhibiting properties of 4-formyl-phenyl boronic acid is at least three times better than those of acetamidophenyl boronic acid.

5 EXAMPLE 3

Storage Stability Test in Liquid Detergent

Phenyl boronic acid derivatives were also tested in storage stability tests in liquid detergents using the method 10 described previously under the following conditions:

Detergent base (US-type)

	% wt (as pure components)
Nansa 1169/p	10.3
15 (Linear Alkylbenzene Sulfonate, LAS)	
Berol 452	3.5
(Alkyl Ether Sulfate, AES)	
Oleic acid	0.5
Coconut fatty acid	0.5
20 Dobanol 25-7	6.4
(Alcohol Ethoxylate, AEO)	
Sodium xylene sulfonate	5.1
Ethanol	0.7
MPG	2.7
25 (Mono Propylene Glycol)	
Glycerol	0.5
Sodium sulfate	0.4
Sodium carbonate	2.7
Sodium citrate	4.4
30 Citric acid	1.5
Water	60.8

Enzyme dosage: 1% w/w Savinase (14 KNPU/g)

Enzyme Stabilizer Dosage: 5 mmole/kg

35 (for boric acid 160 mmole/kg)

22

Storage:

0, 3, 7 and 14 days at 30°C

The results of the inhibition effectiveness IF_i of the phenyl boronic acid enzyme stabilizers tested are listed 5 below:

Inhibitor:	Improvement Factor IF_i
10 -----	-----
Boric acid	1
4-formyl-phenyl-boronic acid	1000
-----	-----

15 For comparison reasons acetamidophenyl boronic acid, 2-formyl-phenyl-boronic acid and 3-formyl-phenyl-boronic acid (all bought at Lancaster) were tested in the same system giving the following results:

20 Inhibitor:	Improvement Factor IF_i
-----	-----
Boric acid	1
acetamidophenyl boronic acid	300
25 2-formyl-phenyl-boronic acid	36
3-formyl-phenyl-boronic acid	230
-----	-----

It appears from the results given above that the storage 30 stability properties of 4-formyl-phenyl boronic acid is at least three times better than those of acetamidophenyl boronic acid, and at least four times better than those of 3-formyl-phenyl-boronic acid, and at least 25 times better than those of 2-formyl-phenyl-boronic acid (all calculated on molar 35 basis).

EXAMPLE 4Storage Stability Test in a Commercial Detergent

5

The inhibition effectiveness IF_i of 4-formyl-phenyl-boronic acid was also found in a commercial detergent Omo Micro.

Omo Micro was bought in a Danish supermarket. The enzymes 10 were inactivated at 90°C (overnight).

The following dosages in the detergent were used:

4-Formyl-phenyl-boronic acid: 1.33 mM, or
Boric acid: 160 mM, and
15 Protease: 1% w/w Savinase (8 KNU/g), and
Lipase: 1% w/w Lipolase (100 KLU/g).

Storage: 0, 7, 15, and 21 days at 40°C.

20 Result: IF_i = 2500.

EXAMPLE 5Storage Stability Test of 4-Carboxybenzeneboronic Acid in
25 Liquid Detergent

4-Carboxybenzeneboronic acid (bought at Lancaster) was tested in a storage stability test in a liquid detergent using the method described previously under the following conditions:

30

Detergent base (US-type)

	% wt (as pure components)
Nansa 1169/p	10.3
(Linear Alkylbenzene Sulfonate, LAS)	
35 Berol 452	3.5

24

(Alkyl Ether Sulfate, AES)

Oleic acid	0.5
Coconut fatty acid	0.5
Dobanol 25-7	6.4

5 (Alcohol Ethoxylate, AEO)

Sodium xylene sulfonate	5.1
Ethanol	0.7
MPG	2.7

(Mono Propylene Glycol)

10 Glycerol	0.5
Sodium sulfate	0.4
Sodium carbonate	2.7
Sodium citrate	4.4
Citric acid	1.5
15 Water	60.8

Enzyme dosage: 1% w/w Savinase (14 KNPu/g)

Enzyme Stabilizer Dosage: 5 mmole/kg


(for boric acid 160 mmole/kg)

20 Storage: 0, 2, 7 and 14 days at 30°C

Result: $IF_1 = 22$.

CLAIMS

1. A liquid composition comprising an enzyme and a phenyl boronic acid derivative enzyme stabilizer of the following formula:



10

where R is selected from the group consisting of hydrogen, hydroxy, C₁-C₆ alkyl substituted C₁-C₆ alkyl, C₁-C₆ alkenyl and substituted C₁-C₆ alkenyl.

15

2. A liquid composition according to claim 1, wherein R is C₁-C₆ alkyl.
3. A liquid composition according to claim 1, wherein R is hydrogen.
4. A liquid composition according to any of claims 1-3, wherein the enzyme is a protease.
5. A liquid composition according to claim 1, additionally comprising a second enzyme, in particular an amylase, a lipase, a cellulase or an oxidoreductase, or any mixture thereof.
6. A liquid composition according to claim 5, wherein the second enzyme is a lipase.


7. A liquid composition according to any of claims 1-6, wherein said phenyl boronic acid derivative enzyme stabilizer is the alkali metal salt of the boronic acid.

5 8. A liquid composition according to any of claims 1-7, wherein said phenyl boronic acid derivative enzyme stabilizer is added in an amount of up to 500 mM, preferably in an amount of 0.001-250 mM, more preferably in an amount of 0.005-100 mM, most preferably in an amount of 0.01-10 mM.

10

9. A liquid detergent composition comprising a surfactant, an enzyme and a phenyl boronic acid derivative enzyme stabilizer of the following formula:

15



20 where R is selected from the group consisting of hydrogen, hydroxy, C₁-C₆ alkyl, substituted C₁-C₆ alkyl, C₁-C₆ alkenyl and substituted C₁-C₆ alkenyl.

10. A liquid detergent composition according to claim 9, 25 wherein R is C₁-C₆ alkyl.

11. A liquid detergent composition according to claim 9, wherein R is hydrogen.

30 12. A liquid detergent composition according to any of claims 9-11, wherein the enzyme is a protease.

13. A liquid detergent composition according to claim 9, additionally comprising a second detergent-compatible enzyme,

27

in particular an amylase, a lipase, a cellulase or an oxidoreductase, or any mixture thereof.

14. A liquid detergent composition according to claim 13,
5 wherein the second enzyme is a lipase.

15. A liquid detergent composition according to any of claims
9-14, wherein said phenyl boronic acid derivative enzyme
stabilizer is the alkali metal salt of the boronic acid.

10

16. A liquid detergent composition according to any of claims
9-15, wherein said phenyl boronic acid derivative enzyme
stabilizer is added in an amount of up to 500 mM, preferably
in an amount of 0.001-250 mM, more preferably in an amount of
15 0.005-100 mM, most preferably in an amount of 0.01-10 mM.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/DK 96/00252

A. CLASSIFICATION OF SUBJECT MATTER

IPC6: C11D 3/386, C11D 3/16

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC6: C11D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

WPI, IFIPAT, CA

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	Biochem.J., Volume 209, No 1, 1983, T. Beesley et al, "The Inhibition of Class C.beta.-lactamases by Boronic Acids" page 229 - page 233	1-3
Y	--	1-16
Y	WO 9512655 A (THE PROCTER AND GAMBLE COMPANY), 11 May 1995 (11.05.95)	1-16
Y	WO 9219707 A (THE PROCTER AND GAMBLE COMAPNY), 12 November 1992 (12.11.92)	1-16
	--	

☒ Further documents are listed in the continuation of Box C. ☒ See patent family annex.

- * Special categories of cited documents
- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed
- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- "Z" document member of the same patent family

Date of the actual completion of the international search

20 Sept 1996

Date of mailing of the international search report

23 -09- 1996

Name and mailing address of the ISA/

Swedish Patent Office

Box 5055, S-102 42 STOCKHOLM

Facsimile No. +46 8 666 02 86

Authorized officer

Teddy Ercégovic

Telephone No. +46 8 782 25 00

Form PCT/ISA/210 (second sheet) (July 1992)

INTERNATIONAL SEARCH REPORT

International application No.

PCT/DK 96/00252

C (Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EP 0478050 A1 (UNILEVER NV), 1 April 1992 (01.04.92)	1-16
A	Biochem.Biophys.Res.Com., Volume 176, No 1, 1991, T.H. Keller et al, "Probing the Specificity of the S1-binding Site of Subtilisin Carlsberg with Boronic Acids" page 401 - page 405	1-16

Form PCT/ISA/210 (continuation of second sheet) (July 1992)

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

05/09/96

PCT/DK 96/00252

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A- 9512655	11/05/95	AU-A- 8095294 EP-A- 0726936 HU-D- 9601188 US-A- 5431842	23/05/95 21/08/96 00/00/00 11/07/95
WO-A- 9219707	12/11/92	AU-A- 2014892 BR-A- 9205958 CA-A- 2109526 CN-B- 1031589 CN-A- 1067265 CZ-A- 9302305 DE-D, T- 69206795 EP-A, B- 0583383 HU-A- 65823 HU-D- 9303086 JP-T- 6507198 NZ-A- 242537 SK-A- 120993 TR-A- 26055 US-A- 5472628	21/12/92 27/09/94 31/10/92 17/04/96 23/12/92 13/04/94 05/09/96 23/02/94 28/07/94 00/00/00 11/08/94 27/06/95 10/08/94 15/12/94 05/12/95
EP-A1- 0478050	01/04/92	AU-A- 8465791 CA-A- 2052077 JP-A- 4283298	09/04/92 25/03/92 08/10/92

Form PCT/ISA/210 (patent family annex) (July 1992)

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.